

# Country Profile

## South Sudan



### *Giraffe Conservation Status Report*

---

#### **Sub-region: East Africa**

*N.B. Although the focus of this profile is on the Republic of South Sudan, reference is made to the historical occurrence of giraffe in Sudan.*

#### **General statistics**

Size of country: 644,329 km<sup>2</sup>

Size of protected areas / percentage protected area coverage: 11.1%

#### **(Sub)species**

Nubian giraffe (*Giraffa camelopardalis camelopardalis*) – possibly

Kordofan giraffe (*G. c. antiquorum*) – possibly

Rothschild's giraffe (*G. c. rothschildi*) – possibly

#### **Conservation Status**

##### IUCN Red List (IUCN 2012):

*Giraffa camelopardalis* (as a species) – least concern

*Giraffa camelopardalis rothschildi* – endangered

*Giraffa camelopardalis camelopardalis* – not assessed

*Giraffa camelopardalis antiquorum* – not assessed

##### In South Sudan:

Under Chapter 5, Section 25 of the Wild Life Conservation and National Parks Act of 2003, no person shall hunt or capture any animal listed in Schedule 1 of the act, which include giraffe.

#### **Issues/threats**

Sudan descended into civil war in 1983. In 2005, after 22 years of war between the National Congress Party (NCP) in the north and the Sudanese People's Liberation Army (SPLA) in the south, the parties signed the Comprehensive Peace Agreement (CPA), putting an end to Africa's longest running conflict. In 2011, the Republic of South Sudan officially became an independent nation.

The armed conflict has severely impacted the lives of communities in and around protected areas in South Sudan, and as such, has resulted in a major assault on the country's wildlife and their habitats (Fay *et al.* 2007; UNEP 2007). Hunting played an important role in human survival during the war and, as a result, uncontrolled and unsustainable hunting decimated wildlife populations (Abdel Magid & Badi 2008; Fay *et al.* 2007; UNEP 2007). Effective management of the country's protected areas throughout the period of the civil war was essentially non-existent while the human, physical, institutional and systemic infrastructure was largely destroyed (Fay *et al.* 2007; UNEP 2007). Moreover, most of these protected areas, while created on paper only, never underwent a consultation process with local stakeholders and protected area boundaries were never demarcated. Additionally, constraints such as inadequate enabling policies and a lack of institutional capacity for wildlife management at operational and administrative levels were a real concern (WCS 2012; UNEP 2007).

With improved stability and prospects for peace and development, the pressure on land and natural resources will rapidly increase in South Sudan (Fay *et al.* 2007; HCENR 2009). Threats to wildlife populations are probably even greater now than during the civil war (Fay *et al.* 2007). Resource extraction plans have begun shortly after the signing of the CPA. Oil exploration concessions have been granted in a large section of the Boma-Jonglei landscape; unsustainable commercial hunting is on the rise as safari hunters are vying for concessions; widespread use of automatic weapons by unauthorized individuals for illegal hunting is common; natural resources have been subject to heavy over-exploitation for agriculture and felling for fuel; and, major infrastructure developments, aided by the international community, are taking place in and around protected areas and important wildlife habitats with little or no environmental assessment, management or planning (WCS 2012; HCENR 2009; Abdel Magid & Badi 2008; Fay *et al.* 2007). Pressures on natural resources continue to increase further due to the inadequately coordinated return of internally displaced people as well as refugees from other countries (HCENR 2009; Fay *et al.* 2007). The effects of these developments will continue to increase pressures on natural resources and competition for land and water, which in turn can have a detrimental effect on wildlife populations (Fay *et al.* 2007).

Data on wildlife numbers for South Sudan is limited. Due to the conflict, little natural resource data has been collected between 1983 and 2005. While recent aerial censuses are helping to fill data gaps, comprehensive population censuses are costly, labour-intensive, and time consuming and it is therefore unlikely that South Sudan's dearth of data will be remedied soon (IRG 2007). Although these aerial surveys indicate that much of South Sudan's terrestrial biodiversity is intact, this does not lessen the existing threats. Critical wildlife species are still at risk, and uncontrolled and unsustainable hunting has caused the local eradication of, among other species, giraffe in certain areas in South Sudan (Fay *et al.* 2007; IRG 2007).

## Estimate population abundance and trends

### Historic

Faunal remains from the Holocene in Central and Eastern Sudan indicate the occurrence of giraffe in the Central Nile valley, Central Nile hinterland, as well as on the Atbara floodplains (Nicoll 2004; Peters 1992). According to East (1999) giraffe formerly occurred widely in southern, central and north-eastern Sudan, with Kordofan giraffe (*G. c. antiquorum*) said to occur to the west of the Nile and Nubian/Rothschild's giraffe (*G. c. camelopardalis/G. c. rothschildi*) to the east. Mackenzie (1954) and Dagg (1962) both documented the general occurrence of giraffe in Sudan. They suggested that Nubian giraffe were fairly common in arid acacia forests and savanna in eastern Sudan bordering Ethiopia; Kordofan giraffe were

scarce in acacia forests and savanna in western and central Sudan; Rothschild's giraffe<sup>1</sup> were fairly common in thin forests of southern Sudan; and Toposa giraffe (*G. c. cottoni*<sup>2</sup>) were common on the plains and foothills east of Kapoeta in Eastern Equatoria (Dagg 1962; Mackenzie 1954).

Several anecdotal records documented the occurrence of giraffe in Sudan from the late 1700s up to the late 1970s, when the first aerial surveys were conducted. Giraffe were apparently widespread and numerous in Southern Darfur and were also documented to occur in Northern Darfur (Wilson 1979 & 1980; Burckhardt 1819; Browne 1799). The species was present in the extreme south-west of the country in the area of Kafia Kingi, and Hofret-en-Nahas to the north-west in the Tebella area in the early 1900s (Wilson 1979). Brocklehurst (1931) suggested that Kordofan giraffe were common throughout the Sudan and extended as far north as the Wadi Howar in Northern Darfur. He documented a herd of several hundred giraffe occurring close to the mouth of the Bahr-el-Ghazal and also noted the occurrence of Rothschild's giraffe in the Mongalla Province in South Sudan. Giraffe were hunted on horseback with swords in southern Kordofan and on the frontier of Abyssinia (today Ethiopia) on the Dinder River (Brocklehurst 1931). In 1927 giraffe were considerably depleted in numbers in the Kordofan Region by a form of rinderpest (Brocklehurst 1931).

Giraffe were seen north of the Wadi Howar in 1932 (Shaw 1936) and were supposedly fairly common in the Geneina and as far north as El Fasher in the late 1940s and early 1950s (Wilson 1980). The species were also recorded in the Kebkabiya area in 1950 (Wilson 1980). Reid (1952) documented a small herd of giraffe occurring on the in the Yei-Juba border and suggested that the species also occurred near Jambo in the Moru District, in the area North of Amadi and near Ngwadi. Sutcliffe (1952) reported giraffe to occur in the Aliab Valley. In Western Darfur, on the borders with Chad, a few giraffe, estimated at 60 animals, were present in 1976 (Wilson 1979). Aerial surveys in the same year indicated a total number of giraffe in Darfur of up to 1,800 animals (Wilson 1979).

The Kordofan giraffe was reportedly scarce by the 1970s (East 1999). Molloy (1951) and Wilson (1979) suggested that giraffe were possibly seriously depleted in numbers owing to the practice of hunting them on horseback with swords or spears. East (1999) indicated that giraffe have been hunted to extinction over most or even all of its former range in the central and north-eastern regions of Sudan. The last herd of giraffe in Dinder National Park in the east was destroyed by illegal hunters in 1985. Giraffe also no longer survived in Radom National Park in the west of Sudan (East 1999).

The first aerial surveys of the Boma National Park (although not legally created at the time due to the disruption of the civil war) in South Sudan were conducted in 1979 to 1980 and estimated the occurrence of some 9,028 animals, supposedly Nubian giraffe (Fryxell 1980a, 1980b). In 1981, an aerial survey of the Southern National Park area estimated 1,325 giraffe inside the park and 770 giraffe in the surrounding area (Boitani 1981). Reconnaissance flights of the Shambe area revealed estimates of some 3,429 animals, also thought to be Nubian giraffe (Hillman *et al.* 1981). Three systematic surveys of the Jonglei area in 1981 and 1982 indicated an estimated 6,025 giraffe (Cobb 1983).

---

<sup>1</sup> Rothschild's giraffe is referred to as Mongalia giraffe by Mackenzie (1954).

<sup>2</sup> *G. c. cottoni* is now considered to be a synonym of *G. c. rothschildi* (Dagg 1971; Seymour 202).

## Recent

Although still present in South Sudan, there is no recent information on the continued occurrence of giraffe in Sudan and the species has possibly become locally extinct. During 22 years of civil war hardly any wildlife surveys were carried out (Fay *et al.* 2007). Ground surveys of the wildlife populations in Boma National Park in 2001 and 2002 confirmed the presence of giraffe (Marjan *et al.* 2002). At the time, aerial surveys could not be conducted due to the ongoing war (WCS 2012).

## Current

The first systematic aerial assessment of wildlife in South Sudan in 25 years was conducted in 2007 by the Wildlife Conservation Society in partnership with local government authorities (Fay *et al.* 2007). The survey covered the most important protected areas and species ranges as well as potential protected areas, including Boma and Southern National Parks as well as the Jonglei area (Fay *et al.* 2007). A population of 404 giraffe were estimated to occur in Boma National Park. Additionally, giraffe were observed during reconnaissance flights over Bandingilo Reserve and viable populations may possibly remain there (Fay *et al.* 2007). However, no giraffe were sighted during aerial surveys of the Jonglei area and the Southern National Park (Fay *et al.* 2007).

Based on the surveys above, the current giraffe numbers for South Sudan are estimated at <450 individuals. However, uncertainty remains as to whether the population(s) are in fact Nubian giraffe or other (sub)species.

## **Future Conservation Management**

The following are proposed conservation management options for giraffe in South Sudan:

- Greater understanding of giraffe population numbers, range and conservation status across the country, including (sub)speciation;
- Development of National Giraffe Strategy for South Sudan;
- Identification of priority conservation efforts for giraffe conservation, specifically for viable remaining populations such as Boma National Park;
- Support to dedicated giraffe conservation, habitat protection, education and awareness initiatives (government, NGO and academic); and
- Assess transboundary giraffe conservation and management initiatives with neighbouring counties i.e. Ethiopia, Uganda and Kenya.

## **Acknowledgements**

This study was financially supported by the Giraffe Conservation Foundation, the Mohamed bin Zayed Species Conservation Fund and Blank Park Zoo.

## References

- Abdel Magid, T.D. & Badi, K.H. 2008. *Ecological zones of the Sudan*. Nile Transboundary Environmental Action Project. Paper presented in the workshop on 'The importance of Wetlands in Sudan'. Khartoum, Sudan.
- Boitani, L. 1981. *The Southern National Park: A Master Plan*. Ministry of Wildlife Conservation and Tourism, Southern Region, Juba-Sudan and Dipartimento per la Cooperazione allo Sviluppo, Ministero degli Affari Esteri. Rome, Italy.
- Brocklehurst, H.C. 1931. *Game animals of the Sudan: their habitats and distribution*. Gurney and Jackson, London, UK.
- Browne, W.G. 1799. *Travels in Africa, Egypt and Syria from the year 1792-1798*. John Murray, London, UK.
- Burckhardt, J.L. 1819. *Travels in Nubia*. John Murray, London, UK.
- Cobb, S.M. 1983. *Government of the Democratic Republic of the Sudan, Ministry of Finance and Economic Planning, Executive Organ of the National Council for Development of the Jonglei canal area*. Technical assistance contract for range ecology survey, livestock investigations and water supply. Final report. Vol. 5. Wildlife studies. Mefit-Babie Srl. Glasgow, Khartoum and Rome.
- Dagg, A.I. 1962. *The distribution of the giraffe in Africa*. School of Graduate Studies, University of Waterloo, Waterloo, Ontario, Canada.
- Dagg, A.I. 1971. *Giraffa camelopardalis*. *Mammalian Species* 5: 1-8.
- East, R. 1999. *African Antelope Database 1998*. IUCN/SSC Antelope Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.
- Fay, M., Elkan, P., Marjan, M. & Grossmann, F. 2007. *Aerial surveys of wildlife, livestock and human activity in and around existing and proposed protected areas of Southern Sudan, dry season 2007. Phase 1*. Wildlife Conservation Society, USA.
- Fryxell, J. 1980a. *Preliminary report on an aerial survey of the Boma National Park region, April 1980*. New York Zoological Society, USA.
- Fryxell, J. 1980b. *Preliminary report on an aerial survey of the Boma National Park region, October 1980*. New York Zoological Society, USA.
- Hillman, A.K.K., Snyder, P.M., Tear, T. & Sommerlatte, M. 1981. *An aerial reconnaissance of the Shambe area, Southern Sudan, 22-26 April 1981*. Report to IUCN/WWF and the Ministry of Wildlife Conservation and Tourism, Southern Region, Sudan. Sudan Open Archive.
- HCENR 2009. *Sudan's fourth national report to the Convention of Biological Diversity*. The Higher Council for Environment and Natural Resources, Khartoum, Sudan.
- IRG. 2007. *Southern Sudan environmental threats and opportunities assessment: Biodiversity and tropical forest assessment*. Prepared by the International Resource Group for the USAID. Washington, DC, USA.
- IUCN 2012. *The IUCN Red List of Threatened Species. Version 2012.1*. <http://www.iucnredlist.org>. Downloaded on 24 August 2012.
- Mackenzie, P.Z. 1954. *Catalogue of wild mammals of the Sudan occurring in the orders Artiodactyla and Perissodactyla*. Sudan Museum (Natural History) Publication No. 4. Khartoum, Sudan.

- Marjan, M.D., Ojok, L.I, Bojo, B.B. & Solomon, I.S. 2002. *The impact of conflict on the Boma National Park: The status of wildlife*. Vol 2: Dry season 2002 wildlife counting. Unpublished manuscript of the Wildlife Survey Team, SACB, USAID, Nairobi, Kenya.
- Molloy, P.G. 1951. Game preservation in the Sudan. *Sudan Wild Life and Sport* **2(1)**: 29-33.
- Nicoll, K. 2004. Recent environmental change and prehistoric human activity in Egypt and Northern Sudan. *Quaternary Science Reviews* **23**: 561-580.
- Peters, J. 1992. *Late Quaternary mammalian remains from Central and Eastern Sudan and their palaeoenvironmental significance*. In Heine, K. (Ed.) Palaeoecology of Africa and the surrounding Islands Vol. 23: 91-115. A.A. Balkema/Rotterdam/Brookfield.
- Reid, R.T.M. 1952. Game notes from Yei and Moru Districts. *Sudan Wild Life and Sport* **2(3)**: 29-39.
- Seymour, R. 2002. *Patterns of Subspecies Diversity in the giraffe, Giraffa camelopardalis (L. 1758): Comparison of Systematic Methods and their implications for Conservation Policy*. PhD Thesis. Institute of Zoology, Zoological Society of London and the Durrell Institute for Conservation and Ecology, University of Kent, UK.
- Shaw, W.B.K. 1936. An expedition in the southern Libyan desert. *Geogr. J.* **87**:193-221.
- Sutcliffe, J.V. 1952. The Game of the Aliab Valley. *Sudan Wild Life and Sport* **2(3)**: 19-22.
- UNEP. 2007. *Sudan: Post-conflict environmental assessment. Synthesis Report*. United Nations Environment Programme, post conflict branch, Nairobi, Kenya.
- Wild Life Conservation and National Parks Act, 2003.
- Wilson, R.T. 1979. Wildlife in southern Darfur, Sudan: Distribution and status at present and in the recent past. *Mammalia* **43(3)**: 323-338.
- Wilson, R.T. 1980. Wildlife in Northern Darfur, Sudan: A review of its distribution and status in the recent past and at present. *Biological Conservation* **17**: 85-101.
- WCS. 2012. *Massive herds found to still exist in Southern Sudan*. Wildlife Conservation Society, USA. <http://www.wcs.org/where-we-work/africa/southern-sudan.aspx> (Accessed 6 June 2012).

### Citation

Marais, A.J., Fennessy, S. & Fennessy, J. 2012. *Country Profile: A rapid assessment of the giraffe conservation status in South Sudan*. Giraffe Conservation Foundation, Windhoek, Namibia.

Map

